

NOAA's 2nd-generation reforecast data set for the NCEP Global Ensemble Forecast System

Tom Hamill

NOAA ESRL, Physical Sciences Division
tom.hamill@noaa.gov

Reforecast background

- 1st-gen reforecast and CFSR have demonstrated utility of reforecasts for weather-climate applications (see previous GMB talks).
- A GEFS reforecast would be a nice complement to CFSR, especially to help with shorter-lead forecasts.
- Were such a data set available, it would have other uses:
 - Spur research in hydrologic prediction, calibration, statistical post-processing methods.
 - Facilitate labs and universities to work with GEFS, improving R2O to EMC.
 - Help diagnose GEFS model error characteristics, especially for low-frequency phenomena, possibly feeding back into GFS development.
 - Help EMC decide on its long-term strategy for reforecasting. How many members, how often?

GEFS reforecast v2 details

- Seeks to mimic GEFS operational configuration as of February 2012.
- Each 00Z, 11-member forecast, 1 control + 10 perturbed.\
- Reforecasts produced every day, for 1984120100 to current (actually, working on 2011 and 2012 now).
- CFSR initial conditions (GSI) + ETR perturbations (cycled with 10 perturbed members). After \sim 22 May 2012, initial conditions from hybrid EnKF/GSI.
- Resolution: T254L42 to day 8, T190L42 from days 7.5 to day 16.
- Fast data archive at ESRL of 99 variables, 28 of which stored at original ~1/2-degree resolution during week 1. All stored at 1 degree. Also: mean and spread to be stored.
- Full archive at DOE/Lawrence Berkeley Lab, where data set was created under DOE grant.
- Web interfaces to each coming soon. Links from TIGGE download pages?

Status

- 00Z reforecasts 1985-2010 completed.
- 2011-12 reforecasts mostly computed, but not yet downloaded to ESRL/PSD and reprocessed.
- Dick Wobus has submitted change request to NCO to save extra GEFS data so we can start archiving real-time data stream.
- We are a few weeks away from opening our ftp/ web sites to public access for 1985-2010:
 - NOAA/ESRL site: fast access, limited data (99 fields).
 - US Department of Energy: slow access, but full data set

Data to be readily available from ESRL

Table 1: Reforecast variables available for selected mandatory and other vertical levels. Φ <u>indicates geopotential</u> height, and an X indicates that this variable is available from the reforecast data set at 1-degree resolution; a Y indicates that the variable is available at the native \sim 0.5 degree resolution. AGL indicates "above ground level."

Vertical	U	V	T	Φ	g	Wind
Level						Power
10 hPa	X	X	X	X		
50 hPa	X	X	X	X		
100 hPa	X	X	X	X		
200 hPa	X	X	X	X		
250 hPa	X	X	X	X		
300 hPa	X	X	X	X	X	
500 hPa	X	X	X	X	X	
700 hPa	X	X	X	X	X	
850 hPa	X	X	X	X	X	
925 hPa	X	X	X	X	X	
1000 hPa	X	X	X	X	X	
$\sigma \approx 0.996$	X	X		X		
$\sigma \approx 0.987$	X	X		X		
σ≅ 0.977	X	X		X		
σ≅ 0.965	X	X		X		
80m AGL	X <u>,Y</u>	X <u>,Y</u>				X,Y


Also: hurricane track files

Data to be readily available from ESRL

Table 2: Single-level reforecast variables archived (and their units). Where $\underline{an}[Y]$ is displayed, this indicates that this variable is available at the native ~ 0.5 -degree resolution as well as the 1-degree resolution.

Variable (units)
Mean sea-level pressure (Pa) [Y]
Skin temperature (K) [Y]
Soil temperature, 0.0 to 0.1 m depth (K) [Y]
Volumetric soil moisture content 0.0 to 0.1 m depth (fraction between
wilting and saturation) [Y]
Water equivalent of accumulated snow depth (kg m ⁻² , i.e., mm) [Y]
2-meter temperature (K) [Y]
2-meter specific humidity (kg kg ⁻¹ dry air) [Y]
Maximum temperature (K) [Y]
Minimum temperature (K) [Y]
10-m u wind component (ms ⁻¹) [Y]
10-m v wind component (ms ⁻¹) [Y]
Total precipitation (kg m ⁻² , i.e., mm) [Y]
Water runoff (kg m ⁻² , i.e., mm) [Y]
Average surface latent heat net flux (W m-2) [Y]
Average sensible heat net flux (W m-2) [Y]
Average ground heat net flux (W m ⁻²) [Y]
Sunshine
Convective available potential energy (J kg-1) [Y]
Convective inhibition (J kg ⁻¹) [Y]
Precipitable water (kg m-2, i.e., mm) [Y]
Total-column integrated condensate (kg m ⁻² , i.e., mm) [Y]
Total cloud cover (%)
Downward short-wave radiation flux at the surface (W m-2) [Y]
Downward long-wave radiation flux at the surface (W m-2) [Y]
Upward short-wave radiation flux at the surface (W m-2) [Y]
Upward long-wave radiation flux at the surface (W m-2) [Y]
Potential vorticity on θ = 320K isentropic surface (K m ² kg ⁻¹ s ⁻¹)
U component on 2 PVU (1 PVU = 1 K m^2 kg ⁻¹ s ⁻¹) isentropic surface (ms ⁻¹)
V component on 2 PVU isentropic surface (ms-1)
Temperature on 2 PVU isentropic surface
Pressure on 2 PVU isentropic surface

Prototype of our web form (creates netCDF4)



Download Reforecast 2 Ensemble Data

This page will will allow you to download one variable at a time. Please note that the more data you request, the slower your response time. Requesting many members and many months/years of data may take several hours to days to process. **Please submit only one request at a time.** If you encounter problems downloading data, please contact esrl.psd.reforecast2@noaa.gov

Select Desired Variables and Associated Levels:

Single Level (1°x1°)	Pressure Levels (1°x1°)	Hybrid Levels (1°x1°) Single Level (Gaussian)
U-Componen U-Componen Convective A Surface Down Surface Upwa Ground Heat Surface Press Volumetric So Sunshine Dun Total Column Maximum Ten Soil Tempera Water Runoff Wind Mixing I Temperature U-Componen	ible Heat Net Flux sure bil Moisture Content ration -Integrated Condensate mperature ture (0-10 cm below surface)	Surface Upward Short-Wave Radiation F Surface Latent Heat Net Flux Mean Sea Level Pressure Precipitable Water Specific Humidity at 2 meters Total Cloud Cover Skin Temperature Minimum Temperature

Select Desired Dates (Available from Dec 1 1984 to Dec 31 2010):

_	 _	maa
From	 То	Ш

- Download all the forecasts within the chosen time period. Help
- ODownload forecasts within the month-days range for the chosen years. Help

Select Desir	ed Dates (Available t	rom Dec 1	1984 to De	ec 31 2010)):				
From		То								
DownloadDownload		asts within th		•	-	elp				
Select Desir	ed Foreca	st Hour(s):								
High Resolu	ution: (Clear)									
0 30 60 108 168 Low Resolu 186 246 306 366	3 33 63 114 174 tion: (Clear) 192 252 312 372	6	9 39 69 126 186 204 264 324 384	12 42 72 132 192	□ 15 □ 45 □ 78 □ 138 □ 216 □ 276 □ 336	184884144222282342	21 51 90 150	□ 24 □ 54 □ 96 □ 156 □ 234 □ 294 □ 354	27 57 102 162	
Select Ensemble Members:										
○ Control										
Select Geog	raphical L	ocation:								
N Select region Point: Latitude:	W E									
Email Addre	ess to Noti	ty When Fi	ie is Ready	/ :	Ret	rieve Data				

Download Reforecast 2 Ensemble Data

This page will will allow you to download one variable at a time. Please note that the more data you request, the slower your response time. Requesting many members and many months/years of data may take several hours to days to process. **Please submit only one request at a time.** If you encounter problems downloading data, please contact esrl.psd.reforecast2@noaa.gov

Variable Levels (1°x1°) Pressure Levels (1°x1°) Hybrid Levels (1°x1°) Single Level (Gaussian)	Sele	ct Desire	ed Variable	s and Ass	sociated Le	evels:						
Variable												
Geopotential Height 1000 925 850 700 500 300 250 200 100 50 10 Temperature U-Component of Wind V-Component of Wind Specific Humidity Temperature U-Component of Wind Specific Humidity To Image: Specific		Single Le	evel (1°x1°)	Pressure	Levels (1º	x1°) Hybr	rid Levels (1º	x1°) Singl	e Level (Gau	ıssian)		
Temperature U-Component of Wind V-Component of Wind Specific Humidity Select Desired Dates (Available from Dec 1 1984 to Dec 31 2010): From To Download all the forecasts within the chosen time period, Help Download forecasts within the month-days range for the chosen years. Help Select Desired Forecast Hour(s): High Resolution: (cien) 0 3 6 9 12 15 18 21 24 27 30 33 36 39 42 45 48 51 54 57 30 60 63 66 69 72 78 84 90 96 102 108 114 120 126 132 138 144 150 156 162 168 174 180 186 192 Low Resolution: (cienr) 186 192 198 204 210 216 222 228 234 240 246 252 258 264 270 276 282 288 294 300 306 312 318 324 330 336 342 348 354 360		Varia	ble		Lev	els (hPa)						
From □ To □ □ ⑤ Download all the forecasts within the chosen time period. Help ⑤ Download forecasts within the month-days range for the chosen years. Help Select Desired Forecast Hour(s): High Resolution: (clear) 0 3 6 9 12 15 18 21 24 27 30 33 36 39 42 45 48 51 54 57 6 63 66 69 72 78 84 90 96 102 10 114 120 126 132 138 144 150 156 162 168 174 180 186 192 186 192 198 204 210 216 222 228 234 240 246 252 258 264 270 276 282 288 294 300 306 312 318 324 330 336 342 348 354 360			Temperature U-Compone V-Compone	ent of Wind ent of Wind								
● Download all the forecasts within the chosen time period. Help	Sele	ct Desire	ed Dates (A	vailable f	rom Dec 1	1984 to De	ec 31 2010)):				
High Resolution: (Clear) 0	DD	ownload a	all the forecas orecasts with	sts within th	e chosen tim th-days rang			elp				
0												
186 192 198 204 210 216 222 228 234 240 246 252 258 264 270 276 282 288 294 300 306 312 318 324 330 336 342 348 354 360 366 372 378 384		0 30 60 108	3 33 63 114	3666120	□ 39□ 69□ 126	 42 72 132	□ 45□ 78	□ 48□ 84	□ 51□ 90	□ 54□ 96	□ 57 □ 102	
Select Ensemble Members:		186 246 306	□ 192□ 252□ 312	□ 258□ 318	□ 264□ 324	270	276	282	□ 288	294	□ 300	
	Sele	ct Enser	nble Memb	ers:								

Other ways of accessing reforecast data

- ftp or gftp into our ftp.cdc.noaa.gov server (if you want lots of the original grib files). [soon]
- Pending Zeus grant, we intend to copy grib fields on Zeus mass store. [MDL plans to use]
- Data written out on external disk here within PSD for special customers (EMC/CPC, OHD), shipped back to them.
- DOE portal to get full dump of single day's grib data (next page).



Web Gateway for Global Ensemble Reforecast Data, Version 2

This web page allows users to download selected days of the full model output from the 2nd-generation NOAA Global Ensemble Forecast System Reforecast (GEFS/R). The format of data downloaded from this page is "grib2" format. It is incumbent on the user to be familiar with the use of this data format as we can provide only minimal user support. For more information on grib2 data, please see GRIB2 use at NCEP.

This reforecast mimics the operational ensemble system that the National Weather Service put into operations in February 2012. The control forecast initial conditions were generated from the Climate Forecast System Reanalysis (CFSR). 10 perturbed initial conditions were generated using the ensemble transform with rescaling (ETR; Wei et al. 2008). Model uncertainty was simulated following Hou et al 2008. Forecasts out to 16 days were generated from 00 UTC initial conditions every day from December 1984 through 2010.

We anticipate that these full model fields provided here will be useful, for example, in providing initial and/or lateral boundary conditions for regional reforecasts with various limited-area models. To access a subset of model output, for example a small number particular fields such as precipitation, surface temperatures, etc., please use the interface at ESRL/PSD. For a more complete description of this reforecast data set, please read [insert URL].

Please submit only one request at a time. If you encounter problems downloading data, please contact esrl.psd.reforecast2@noaa.gov

This 2nd-generation GEFS/R was generated under a DOE supercomputer grant at Lawrence Berkeley Lab.


	esired Date (from Dec. 1, Dec. 31, 2010):
Date	##
Select E	nsemble Members:
Control:	Perturbation:
	1 2 3 4 5 6 7 8 9 10
	Select All or Clear
Email Ac	Idress to Notify When File

This DOE site will be ready for use in a few weeks for access to tape storage of full data (slower).

Use this to access full model state.

Later in this presentation we will show a regional WRF reforecast initialized from GEFS reforecast.

500 hPa Z anomaly correlation (from deterministic control)




How does the skill of post-processed forecasts change from reforecast v1 to v2?

 Let's look at post-processed 24-h precipitation forecast guidance.

 Will use a "rank analog" approach for postprocessing (description in supplementary slides)


Skill of calibrated precipitation forecasts

(over US, 1985-2009, "rank analog" calibration method)




Skill of calibrated precipitation forecasts


(over US, 1985-2009, "rank analog" calibration method)




Reliability, > 10 mm precipitation 24 h⁻¹




Reliability, > 50 mm precipitation 24 h⁻¹






The bad and ugly atmospheric rivers case study

February 16, 2004 12-24 UTC




➤ 10" rain in the coastal mountains, 4-7" in Russian River watershed. Streamflows in top 0.2% of historical observations.

(a) 24-h accumulated precip analysis, VT = 2004021700




(c) P(5-6 day accum precip > 10 mm), Reforecast v2, VT = 2004021700


Analyzed precipitation amount (mm)



(b) 5-6 day mean forecast, Reforecast v2, VT =2004021700




(d) P(5-6 day accum precip > 50 mm), Reforecast v2, VT = 2004021700




6-day forecast

(a) 24-h accumulated precip analysis, VT = 2004021700




Analyzed precipitation amount (mm)

(c) P(3-4 day accum precip > 10 mm),Reforecast v2, VT = 2004021700




Probability (3-4 day precip > 10 mm)

(b) 3-4 day mean forecast, Reforecast v2, VT = 2004021700




(d) P(3-4 day accum precip > 50 mm),Reforecast v2, VT = 2004021700



Probability (3-4 day precip > 50 mm)


4-day forecast

(a) 24-h accumulated precip analysis, VT = 2004021700




(c) P(1-2 day accum precip > 10 mm), Reforecast v2, VT = 2004021700

Analyzed precipitation amount (mm)




Probability (1-2 day precip > 10 mm)


(b) 1-2 day mean forecast, Reforecast v2, VT =2004021700



(d) P(1-2 day accum precip > 50 mm), Reforecast v2, VT = 2004021700




2-day forecast




Colors: over the ocean, the total-column precipitable water. Over land, the 24-h accum. precipitation.

Wind barbs for the 925 hPa level.

Red contours: mean sea-level pressure.



4-day forecast



2-day forecast


Case study, tentative conclusions

- Statistical post-processing will not be able to correct for everything. In this case, the synoptic-scale predictability was apparently quite low.
- Improvements to post-processed probabilistic forecasts in such a case will require improved ensemble guidance.

The "good" atmospheric rivers case study: Nov 2006 Oregon-Washington floods


November 07, 2006 00-12 UTC

SSMI Water Vapor (Wentz algorithm)




8-20 inches of rain in Cascades; flooded rivers; extensive damage to Mt. Rainier NP.

(a) 24-h accumulated precip analysis, VT = 2006110700




(b) 5-6 day mean forecast,

Reforecast v2, VT = 2006110700



(c) P(5-6 day accum precip > 10 mm),Reforecast v2, VT = 2006110700

Analyzed precipitation amount (mm)




Probability (5-6 day precip > 10 mm)


(d) P(5-6 day accum precip > 50 mm),Reforecast v2, VT = 2006110700

0 0.1 0.5 1 2.5 5 10 25 50 100 150

Ensemble-mean forecast amount (mm)




(a) 24-h accumulated precip analysis, VT = 2006110700




(c) P(3-4 day accum precip > 10 mm),

Analyzed precipitation amount (mm)




Probability (3-4 day precip > 10 mm)

(b) 3-4 day mean forecast, Reforecast v2, VT =2006110700




(d) P(3-4 day accum precip > 50 mm), Reforecast v2, VT = 2006110700




4-day forecast

(a) 24-h accumulated precip analysis, VT = 2006110700




(c) P(1-2 day accum precip > 10 mm),

Analyzed precipitation amount (mm)




Probability (1-2 day precip > 10 mm)


(b) 1-2 day mean forecast, Reforecast v2, VT = 2006110700



(d) P(1-2 day accum precip > 50 mm),Reforecast v2, VT = 2006110700




2-day forecast




Colors: over the ocean, the total-column precipitable water. Over land, the 24-h accum. precipitation.

Wind barbs for the 925 hPa level.

Red contours: mean sea-level pressure.







Inconsistent forecasts of orientation of isobars and hence wind direction

Spaghetti Westerns

Consistent forecasts of orientation of isobars and hence wind direction



The bad and the ugly



Demo: Regional reforecast with WRF ARW v3.4 using global reforecast for initial, boundary conditions

- 2-way nested simulation 36-, 12- and 4-km with 36 vertical levels
 - 12- and 4-km moving nests
- Time step: 180, 60, and 20 s
- Initial and boundary condition: GFS reforecast ensemble member
- Tiedtke cumulus scheme on 36 and 12 km; explicit on 4 km
- YSU PBL scheme
- HYCOM ocean analysis
- WSM6 microphysics
- Noah land surface
- 2D Smagorinsky turbulence scheme
- Goddard shortwave radiation
- RRTM longwave radiation
- Second order diffusion
- Positive definite scalar advection
- Donelan wind-dependent drag formulation
- Garratt wind-dependent enthalpy surface fluxes


2005 Rita official forecast (Houston, TX evacuated)





TC Rita (2005)

GFS reforecast ensemble

72-h forecast initialized at 00Z 22 Sept










TC Rita (2005)

ARW ensemble with GFS reforecast ensemble as boundary and initial conditions

72-h forecast initialized at 00Z 22 Sept










A synthetic example of using reforecasts to make track error bias corrections

72-h Forecast Verifying 1200 UTC 9 September








Red **f**: mean forecast position

Blue dot: forecast positions of +72-h forecast analogs

End of red tail ____: observed positions at +72 h

Application: diagnosis of MJO



Let's examine subset of cases that start off with initial conditions that project onto relatively strong MJO emerging from African continent.


We have lots of cases of this afforded by multi-decadal reforecasts.

It appears that GEFS is too regular with its RMM1/RMM2 forecasts, while the analyzed evolution is more scattered with its trajectories.

Other work in progress

- Blocking diagnostics (w. Steve Colucci, Cornell U.)
- Development of forecast tools to support extendedrange renewable-energy forecasts

Extreme Forecast Index of 80-m wind speed (+ sea-level pressure) 72-h forecast valid 2007120300




Extreme Forecast Index

Example of ECMWF's Extreme Forecast Index, where the current ensemble guidance is compared to the model forecast climatology. Values near 1.0 indicate that ensemble forecasts are consistently higher than forecasts populating the climatology.

This can be useful especially in situations where there isn't observational data available for calibration.

Application: extended-range tornado forecasting


4/11/1996 Forecast, 204-hour through 276-hour leadtime
Using 3 PCs of 0-6 km Shear, log(CAPE) & Conv.Precip. as Predictors for Logistic Regression
Probability of tornado (>EF0) event



Francisco Alvarez,
St. Louis University,
is working with me
and others on using the
reforecasts to make
extended-range
predictions of
tornado probabilities.

Ph.D. work, in progress.

Application: Improved 6-10 day and week-2 forecast guidance from CPC



Dan Collins, CPC leading this effort.

Conclusions


- Reforecast data set created for current operational NCEP GEFS.
- Reforecast data and real-time forecasts available shortly.
- Useful for statistical calibration.
- Helpful for determining how you may choose to configure real-time reforecasts in the future.

Supplementary slides


Basic analog technique for statistical downscaling (here, v1)

Today's ens. mean forecast + subsequent analyzed precipitation


26 Nov 2005




On the left are old forecasts similar to today's ensemblemean forecast. For making probabilistic forecasts, form an ensemble from the accompanying analyzed weather on the right-hand side.



Analog technique for statistical downscaling



Problem with basic analog reforecast technique



Say today's forecast is for 20 mm. There are more forecasts slightly less than 20 mm than slightly more than 20 mm.

Assuming correlation between forecast and observations, analogs will be biased toward lower precipitation amounts.


"Rank" analog procedure

 Convert precipitation forecast time series to ranks:

```
x = [0, 0, 7, 15, 1, 3, 6, 4, 1, 2, 12, 5, 6, 8]
x(r) = [1.5, 1.5, 11, 14, 3.5, 6, 9, 7, 3.5, 5, 13, 8, 10, 12]
```

"Rank" analog procedure


 Convert precipitation forecast time series to ranks:



with standard analog, these would be the two forecasts with the closest values.

"Rank" analog procedure

 Convert precipitation forecast time series to ranks:



with rank analog, these would be the two forecasts with the closest ranks.

Rank analog calibration details

- 24-h accumulated precipitation, validated on NARR grid (~32 km) over CONUS, 1985-2009.
- Rank analog approach: at each grid point in CONUS, using that grid point and +/- 3 surrounding grid points in N-S, E-W direction, find dates of 75 past forecasts that are closest in average precipitation rank of ensemble mean forecast. Make probabilistic forecasts from analyzed NARR precipitation data on dates of those 75 analogs.
- (Conventionally calculated) Brier Skill Scores, reliability diagrams, etc. NARR again used for verification.


Regional Reforecast with ARW v3.4


- 2-way nested simulation 36-, 12- and 4-km with 36 vertical levels
 - 12- and 4-km moving nests
- Time step: 180, 60, and 20 s
- Initial and boundary condition: GFS reforecast ensemble member
- Tiedtke cumulus scheme on 36 and 12 km; explicit on 4 km
- YSU PBL scheme
- HYCOM ocean analysis
- WSM6 microphysics
- Noah land surface
- 2D Smagorinsky turbulence scheme
- Goddard shortwave radiation
- RRTM longwave radiation
- Second order diffusion
- Positive definite scalar advection
- Donelan wind-dependent drag formulation
- Garratt wind-dependent enthalpy surface fluxes


TC Rita (2005)

GFS reforecast ensemble

72-h forecast initialized at 00Z 22 Sept










TC Rita (2005)

ARW ensemble with GFS reforecast ensemble as boundary and initial conditions

72-h forecast initialized at 00Z 22 Sept









Some known applications in development

- Real-time web page of experimental PQPF forecasts (ESRL, in development to replace v1 web page)
- Development of experimental forecast products for renewable energy sector (Hamill, ESRL).
- 6-10 day and 8-14 day forecasts (Dan Collins, CPC).
- Detecting and correcting bias in hurricane track and intensity (Jiayi Peng, NCEP/EMC, Tom Galarneau, NCAR).
- Objective probabilities for severe weather several days to weeks hence (Francisco Alvarez, St. Louis Univ. + Hamill)
- Improving hydrologic predictions (Haksu Lee et al., NOAA/ NWS/OHD).
- Also: BAMS article once published will presumably spur wider usage.

Scores, > 10 mm and > 50 mm





Brier Skill Scores, Day +0-1, 1.0 mm 24h⁻¹ (reforecast version 2)




Brier Skill Scores, Day +0-1, 1.0 mm 24h⁻¹


(reforecast version 1)








Brier Skill Score




NARR climatology, > 1-mm 24 h⁻¹



a modest relationship between where precipitation is common and the magnitude of skill.

Also, note relationship with terrain forcing.


Brier Skill Scores, Day +0-1, 10.0 mm 24h⁻¹ (reforecast version 2)




Brier Skill Score

Brier Skill Score


Brier Skill Scores, Day +0-1, 10.0 mm 24h⁻¹ (reforecast version 1)



NARR climatology, $> 10 \text{ mm } 24 \text{ h}^{-1}$



Reliability, > 1 mm precipitation 24 h⁻¹



Reliability, > 1 mm precipitation 24 h⁻¹

